

Adaptations for Carbon and Energy Metabolism in *Psychrobacter* 273-4

Peter W. Bergholz

Center for Genomic and Evolutionary Studies on Microbial Life at Low Temperature
Michigan State University
Plant and Soil Science Room 545
East Lansing, MI 48824
USA
bergholz@msu.edu

Hector L. Ayala-del-Rio

Center for Genomic and Evolutionary Studies on Microbial Life at Low Temperature
University of Puerto Rico at Humacao
USA

Natalia N. Ivanova

Microbial Genome Analysis Program
DOE Joint Genome Institute
USA

Michael F. Thomashow

Center for Genomic and Evolutionary Studies on Microbial Life at Low Temperature
Michigan State University
USA

James M. Tiedje

Center for Genomic and Evolutionary Studies on Microbial Life at Low Temperature
Michigan State University
USA

Psychrobacter 273-4 was isolated after 10,000-40,000 yr exposure to continuous subzero temperatures and low water activity. Understanding the genetic and physiological adaptations that *Psychrobacter* has made to inhabit permafrost will enable us to hypothesize about potential microbial life in extraterrestrial cryo environments. In this study, we asked the question, “What carbon sources has *Psychrobacter* adapted to use for optimal growth at low temperatures?” Metabolic pathway analysis indicated that, despite previous expectations, *Psychrobacter* was not adapted to grow on sugars, but appeared to have the capacity to grow on carboxylic acids. Metabolism of the 2-C compound glyoxylate is likely accomplished through an unusual pathway ending in 3-hydroxy-aspartate. Microarray comparisons of *Psychrobacter* growth at 22°C vs 4°C showed that *Psychrobacter* upregulates genes for glycolate uptake and glyoxylate metabolism at 4°C suggesting increased demand for carbon and energy from this substrate during low temperature metabolism. Sequence analysis of *icl* (isocitrate lyase) gene has shown that it clusters with *icl* genes from other cold adapted organisms. These data led to the hypothesis that *Psychrobacter* is adapted for metabolism of carboxylic acids, particularly the 2-C compounds acetate and glyoxylate. *Psychrobacter* was grown in a defined media containing several different compounds. Lactate, pyruvate, and acetate all supported growth as sole carbon sources. Growth rates on acetate were more than thrice the rates

observed in 1/2 tryptic soy broth at 4°C. This work shows *Psychrobacter* is adapted to use oxidized substrates, and suggests the glyoxylate bypass (involving *icl*) plays an important role in low temperature energy metabolism.